

STIMULI-SENSITIVE DRUG DELIVERY SYSTEMS

Vladimir Torchilin

Department of Pharmaceutical Sciences and Center for Pharmaceutical Biotechnology and Nanomedicine, Northeastern University, Boston, MA 02115, USA

Drug carriers are used to protect sensitive drug molecules from the inactivation and to improve drug's pharmacokinetics and delivery to the site of disease. Ideally, drug carrier should: (a) accumulate in required organ or tissue, and (b) penetrate inside target cells delivering there its load. Organ or tissue (tumor, infarct) accumulation could be achieved by the passive targeting via the enhanced permeability and retention effect or by the specific ligand(antibody)-mediated active targeting, while the intracellular delivery could be mediated by certain internalizable ligands (folate, transferrin) or by cell-penetrating peptides. To be able to behave this way, drug carrier should simultaneously carry on its surface various moieties capable of functioning in a certain orchestrated order.

Now, within the frame of this concept, multifunctional stimuli-responsive nanocarriers are developed, i.e. nanocarriers that, depending on the particular requirements, can circulate long; target the site of the disease via both non-specific and/or specific mechanisms; respond local stimuli characteristic of the pathological site by, for example, releasing an entrapped drug or deleting a protective coating facilitating thus the contact between drug-loaded nanocarriers and target cells; and even provide an enhanced intracellular delivery of an entrapped drug with its subsequent delivery to specific intracellular organelles, such as nuclei, lysosomes or mitochondria. Such carriers can be additionally supplemented with reporter moieties to follow their real-time biodistribution and target accumulation. Among new developments to be considered in the area of multifunctional pharmaceutical nanocarriers are: drug- or/and RNA-loaded delivery systems additionally decorated with cell-penetrating peptides for the enhanced intracellular delivery; "smart" multifunctional drug delivery systems, which can reveal/expose temporarily hidden functions under the action of certain local stimuli characteristic for the pathological zone (such as lowered pH, redox-conditions, hypoxia, or locally increased expression of certain enzymes); new means for controlled delivery and release of siRNA; approaches for intracellular drug delivery and organelle targeting; and application of nanocarriers co-loaded with siRNA and drugs to treat multidrug resistant tumors.